

CANADIAN GUIDELINES ON SEXUALLY TRANSMITTED INFECTIONS (2008 UPDATE)

IV – MANAGEMENT AND TREATMENT OF SPECIFIC SYNDROMES

URETHRITIS (January 2008)

References

1. Swartz SL, Kraus SJ, Herrmann KL, Stargel MD, Brown WJ, Allen SD. Diagnosis and etiology of nongonococcal urethritis. *J Infect Dis.* 1978;138:445–454.
2. McKee KT Jr, Jenkins PR, Garner R, et al. Features of urethritis in a cohort of male soldiers. *Clin Infect Dis.* 2000;30:736–741.
3. Wendel KA, Erbeling EJ, Gaydos CA, Rompalo AM. Use of urine polymerase chain reaction to define the prevalence and clinical presentation of *Trichomonas vaginalis* in men attending an STD clinic. *Sex Transm Infect.* 2003;79:151–153.
4. Madeb R, Nativ O, Benilevi D, Feldman PA, Halachmi S, Srugo I. Need for diagnostic screening of herpes simplex virus in patients with nongonococcal urethritis. *Clin Infect Dis.* 2000;30:982–983.
5. Mena L, Wang X, Mroczkowski TF, Martin DH. *Mycoplasma genitalium* infections in asymptomatic men and men with urethritis attending a sexually transmitted diseases clinic in New Orleans. *Clin Infect Dis.* 2002;35:1167–1173.
6. Dupin N, Bijaoui G, Schwarzingler M, et al. Detection and quantification of *Mycoplasma genitalium* in male patients with urethritis. *Clin Infect Dis.* 2003;37:602–605.
7. Bradshaw CS, Denham IM, Fairley CK. Characteristics of adenovirus associated urethritis. *Sex Transm Infect.* 2002;78:445–447.
8. Azariah S, Reid M. Adenovirus and non-gonococcal urethritis. *Int J STD AIDS.* 2000;11:548–550.
9. Varela JA, Otero L, Garcia MJ, et al. Trends in the prevalence of pathogens causing urethritis in Asturias, Spain, 1989–2000. *Sex Transm Dis.* 2003;30:280–283.
10. Grosskurth H, Mayaud P, Mosha F, et al. Asymptomatic gonorrhoea and chlamydial infection in rural Tanzanian men. *BMJ.* 1996;312:277–280.
11. Martin DH, Bowie WR. Management of STD syndromes in men. In: Holmes KK, Sparling P, Mardh PA, et al, eds. *Sexually Transmitted Diseases.* 3rd ed. New York, NY: McGraw Hill; 1999: 833-845.
12. Simmons PD. Evaluation of the early morning smear investigation. *Br J Vener Dis.* 1978;54:128–129.
13. Burstein GR, Zenilman JM. Nongonococcal urethritis — a new paradigm. *Clin Infect Dis.* 1999;28(suppl 1):S66–73.
14. Centers for Disease Control and Prevention (CDC). Screening tests to detect *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infections — 2002. *MMWR Recomm Rep.* 2002;51(RR-15): 1-27.
15. Handsfield HH, McCormack WM, Hook EW 3rd, et al. A comparison of single-dose cefixime with ceftriaxone as treatment for uncomplicated gonorrhoea. The Gonorrhoea Treatment Study Group. *N Engl J Med.* 1991;325: 1337–1341.
16. Stamm WE, Hicks CB, Martin DH, et al. Azithromycin for empirical treatment of the nongonococcal urethritis syndrome in men. A randomized double-blind study. *JAMA.* 1995;274:545–549.
17. Steingrimsson O, Olafsson JH, Thorarinsson H, Ryan RW, Johnson RB, Tilton RC. Single dose azithromycin treatment of gonorrhoea and infections caused by *C. trachomatis* and *U. urealyticum* in men. *Sex Transm Dis.* 1994;21:43–46.

18. Lau CY, Qureshi AK. Azithromycin versus doxycycline for genital chlamydial infections. A meta-analysis of randomized clinical trials. *Sex Transm Dis.* 2002;29:497–502.
19. Macke BA, Maher JE. Partner notification in the United States: an evidence-based review. *Am J Prev Med.* 1999;17:230–242.
20. Public Health Agency of Canada. *Interim Statement on the Treatment of Gonorrhoea In Canada.* Ottawa, ON: Public Health Agency of Canada; November 2004. Available at: www.phac-aspc.gc.ca/std-mts/pdf/is-gonorrhoea-2004_e.pdf. Accessed March 1, 2005.
21. Borchardt KA, al-Haraci S, Maida N. Prevalence of *Trichomonas vaginalis* in a male sexually transmitted disease clinic population by interview, wet mount microscopy and the InPouch TV test. *Genitourin Med.* 1995;71:405–406.
22. Lautenschlager S, Eichmann A. Urethritis: an underestimated clinical variant of genital herpes in men? *J Am Acad Dermatol.* 2002;46:307–308.