

# Foreword

Table of Contents

Expert Working Group on the  
Canadian Guidelines on Sexually  
Transmitted Infections

Acknowledgments

Introduction

Levels and Quality of Evidence for  
Treatment Recommendations

Important Considerations for  
Guidelines Users

# TABLE OF CONTENTS

---

---

## **Section 1**

*Expert Working Group on the  
Canadian Guidelines on Sexually  
Transmitted Infections*

*Acknowledgements*

*Introduction*

*Levels and Quality of Evidence  
for Treatment Recommendations*

*Important Considerations for  
Guidelines Users*

---

## **Section 2**

*Primary Care and  
Sexually Transmitted Infections*

---

## **Section 3**

*Laboratory Diagnosis  
of Sexually Transmitted Infections*

*Collection and Transportation of Specimens*

*Laboratory Testing Methods*

*Laboratory Diagnosis of Specific Infections*

## **Section 4**

---

### ***Management and Treatment of Specific Syndromes***

*Syndromic Management  
of Sexually Transmitted Infections*

*Epididymitis*

*Genital Ulcer Disease (GUD)*

*Pelvic Inflammatory Disease (PID)*

*Prostatitis*

*Sexually Transmitted Intestinal  
and Enteric Infections*

*Urethritis*

*Vaginal Discharge (Bacterial Vaginosis,  
Vulvovaginal Candidiasis, Trichomoniasis)*

## **Section 5**

---

### ***Management and Treatment of Specific Infections***

*Chancroid*

*Chlamydial Infections*

*Ectoparasitic Infestations (Pubic Lice, Scabies)*

*Genital Herpes Simplex Virus (HSV) Infections*

*Genital Human Papillomavirus (HPV) Infections*

*Gonococcal Infections*

*Hepatitis B Virus Infections*

*Human Immunodeficiency Virus (HIV) Infections*

*Lymphogranuloma Venereum (LGV)*

*Syphilis*

---

## **Section 6**

### **Specific Populations**

*Immigrants and Refugees*

*Inmates and Offenders*

*Men Who Have Sex with Men (MSM)/  
Women Who Have Sex with Women (WSW)*

*Pregnancy*

*Sexual Abuse in Peripubertal and  
Prepubertal Children*

*Sexual Assault in Postpubertal  
Adolescents and Adults*

*Sex Workers*

*Substance Use*

*Travellers*

## **Section 7**

---

### **Appendices**

*A: Patient Counselling Guide on Condom Use*

*B: How to Use a Male Condom/  
How to Use a Female Condom*

*C: Resources and Reference Tools  
for Health Professionals*

*D: Provincial and Territorial Offices  
for the Prevention and Control of STI*

*E: Provincial Laboratories*

*F: Forensic Evidence, Services and Laboratories*

*G: Referral Centres for STIs  
in Peripubertal and Prepubertal Children*

*H: Tanner Scale of Sexual Maturity*

## **Section 8**

---

### **General index**

# EXPERT WORKING GROUP FOR THE CANADIAN GUIDELINES ON SEXUALLY TRANSMITTED INFECTIONS

<b>Chair</b>	<b>Tom Wong</b> , MD, MPH, FRCPC, Director, Community Acquired Infections Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada
<b>Section Chairs</b>	<p><b>Primary Care and Sexually Transmitted Infections</b>  <b>Marc Steben</b>, MD, FCFP, médecin-conseil, Direction risques biologiques, environnementaux et occupationnels, Institut national de santé publique du Québec et clinique des maladies de la vulve, Hôpital Notre-Dame, Centre hospitalier de l'Université de Montréal</p> <p><b>Laboratory Diagnosis of Sexually Transmitted Infections</b>  <b>Max Chernesky</b>, PhD, Professor Emeritus, McMaster University</p> <p><b>Management and Treatment of Specific Syndromes</b>  <b>Mark Yudin</b>, MD, MSc, FRCSC, Assistant Professor, University of Toronto; Deputy Head, Department of Obstetrics and Gynecology, St. Michael's Hospital</p> <p><b>Management and Treatment of Specific Infections</b>  <b>Barbara Romanowski</b>, MD, FRCPC, Clinical Professor of Medicine, Division of Infectious Diseases, Faculty of Medicine and Dentistry, University of Alberta</p> <p><b>Specific Populations</b>  <b>Rhonda Kropp</b>, BScN, MPH, Head, Strategic Analysis and Knowledge Transfer, Sexual Health and STI Section, Community Acquired Infections Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada</p>
<b>Members</b>	<p><b>Joanne Embree</b>, MD, FRCPC, Departments of Medical Microbiology &amp; Pediatrics and Child Health, University of Manitoba</p> <p><b>William Fisher</b>, PhD, Professor, Departments of Psychology and Obstetrics and Gynaecology, University of Western Ontario</p> <p><b>Cathy Latham-Carmanico</b>, RN, BScN, Knowledge Development and Research Analyst, Sexual Health and STI Section, Community Acquired Infections Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada</p> <p><b>Lai-King Ng</b>, PhD, Director of Bacteriology and Enteric Diseases Program, National Microbiology Laboratory, Public Health Agency of Canada</p> <p><b>David Patrick</b>, MD, MHSc, FRCPC, Associate Professor, UBC Healthcare &amp; Epidemiology; Director, Epidemiology Services, British Columbia Centre for Disease Control</p> <p><b>Michael Rekart</b>, MD, DTM&amp;H, MHSc, Director, HIV/AIDS Control, British Columbia Centre for Disease Control</p> <p><b>Ameeta Singh</b>, MD, BMBS, MSc, FRCPC, Infectious Diseases Medical Consultant, Alberta Health and Wellness; Clinical Associate Professor, Department of Medicine, University of Alberta; Medical Director, Capital Health STD Centre</p>
<b>Secretariat</b>	<b>Allison Ringrose</b> , BHSc, Prevention and Research Analyst, Sexual Health and STI Section, Community Acquired Infections Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada

## ACKNOWLEDGMENTS

With the assistance of the Sexual Health and Sexually Transmitted Infections Section, Community Acquired Infections Division, Centre for Communicable Diseases & Infection Control, Public Health Agency of Canada:

Editor-in-Chief:	Dr. Tom Wong
Associate Editor:	Cathy Latham-Carmanico
Production Coordinator:	Robert Lerch
Editing and Production Support:	Tory Atwood, Barbara Jones, Leila Khalaf, Rhonda Kropp, Jacques Néron, Allison Ringrose, Dr. Maxim Trubnikov

Many health professionals from across Canada volunteered their time to author chapters for these guidelines, and their participation is acknowledged by chapter:

**Fred Y. Aoki**, MD, Professor of Medicine, Medical Microbiology and Pharmacology & Therapeutics Member, Section of Adult Infectious Diseases, Faculty of Medicine, University of Manitoba, Author: Genital Herpes simplex virus (HSV) Infections; **Dr. Max Chernesky**, PhD, Professor Emeritus, McMaster University, Author: Laboratory Diagnosis of Sexually Transmitted Infections; **Francois Coutlée**, Clinical Researcher, Laboratoire de Virologie Moléculaire, Centre de Recherche Centre Hospitalier de l'université de Montréal Hôpital Notre-Dame, Co-Author : Genital Human Papillomavirus (HPV) Infections; **Laurent Delorme**, MD CSPQ FRCP(C), médecin microbiologiste infectiologue, Hôpital Charles-LeMoine, Co-Author : Genital Ulcer Disease (GUD); **Francisco Diaz-Mitoma**, M.D., Ph.D., FRCPC, Professor and Chief, Division of Virology, Children's Hospital of Eastern Ontario, University of Ottawa, Co-Author : Genital Ulcer Disease (GUD); **Alex Ferenczy**, MD, Professor of Pathology and Obstetrics & Gynecology, McGill University and the Sir Mortimer B. Davis-Jewish General Hospital Montreal, Co-Author : Genital Human Papillomavirus (HPV) Infections; **William A. Fisher**, PhD, Professor, Departments of Psychology and Obstetrics and Gynaecology, University of Western Ontario, Author: Primary Care and Sexually Transmitted Infections; **Sarah Forgie**, MD FRCPC, Assistant Professor, Pediatrics, Division of Infectious Diseases, University of Alberta, Associate Director, Infection Control, Stollery Children's Hospital and University of Alberta Hospital, Co-Author: Sexual Abuse in Peri-Pubertal and Prepubertal Children, Sexual Assault in Post Pubertal Adolescents and Adults; **Eduardo L. Franco**, MPH, DrPH, James McGill Professor of Epidemiology and Oncology Director, Division of Cancer Epidemiology, McGill University, Co-Author : Genital Human Papillomavirus (HPV) Infections; **Deana Funaro**, MD, FRCPC dermatology, Clinical professor at Notre-Dame Hospital and Ste-Justine Hospital, University of Montreal, Co-Author : Genital Ulcer Disease (GUD); **David Haase**, MD, Professor, Department of Medicine, Division of Infectious Diseases, Dalhousie University, Author: Human Immunodeficiency Virus (HIV) Infections; **Rhonda Kropp**, BScN, MPH, Head, Strategic Analysis and Knowledge Transfer, Sexual Health and STI Section, Community Acquired Infections Division, Public Health Agency of Canada, Author: Men Who Have Sex With Men (MSM)/Women Who Have Sex With Women (WSW), Substance Use, Co-Author: Lymphogranuloma Venereum (LGV); **Annie-Claude Labbé**, MD, FRCPC, Department of microbiology, Hôpital Maisonneuve-Rosemont Montreal, Co-Author : Genital Ulcer Disease (GUD); **Cathy Latham-Carmanico**, RN, BScN, Knowledge Development and Research Analyst, Sexual Health and STI Section, Community Acquired Infections Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada, Author: Primary Care and Sexually Transmitted Infections; **Janice Mann**, MD, Senior Knowledge Transfer Advisor, Strategic Communications and Knowledge Exchange, Canadian Agency for Drugs and Technologies in Health, Author: Inmates and Offenders, Lymphogranuloma Venereum (LGV); **Lynette J. Margesson**, MD, FRCPC, Assistant Professor of Obstetrics and Gynecology and of Medicine(Dermatology)Dartmouth Medical School, Co-Author : Genital Ulcer Disease (GUD); **Deborah M. Money**, MD, FRCSC, Associate Professor, University of British Columbia, B.C. Women's Hospital, Co-Author: Pregnancy; **Gina Olgilvie**, MD, MSc, University of British Columbia, Vancouver, Author: Urethritis; **Ron Read**, MD, Head, Infectious Diseases, Department of Medicine, Microbiology and Infectious Diseases, University of Calgary/Calgary Health Region, Author: Hepatitis B Virus Infections; **Michael L. Rekart**, MD, DTM&H, MHSc, Director, HIV/AIDS Control, British Columbia Centre for Disease Control, Author: Sex Workers; **Barbara Romanowski**, MD, FRCPC,

Clinical Professor of Medicine, Division of Infectious Diseases, Faculty of Medicine and Dentistry, University of Alberta, Author: Ectoparasitic Infestations, Gonococcal Infections; **Allan Ronald**, MD, Distinguished Professor Emeritus, University of Manitoba, Author: Chancroid; **Shelly Sarwal**, MD, MSc, FRCP(C), Medical Officer of Health, Nova Scotia Department of Health, Author: Vaginal Discharge; **Rita Shahin**, MD, Toronto Public Health, Author: Travellers; **Ameeta Singh**, BMBS, MSc, FRCPC, Infectious Diseases Medical Consultant, Alberta Health and Wellness Clinical Associate Professor, Department of Medicine, University of Alberta, Medical Director, Capital Health STD Centre, Author: Sexual Abuse in Peri-Pubertal and Prepubertal Children, Sexual Assault in Post Pubertal Adolescents and Adults, Syphilis; **Marc Steben**, MD, médecin-conseil, Direction risques biologiques, environnementaux et occupationnels, Institut national de santé publique du Québec et clinique des maladies de la vulve hôpital Notre-Dame Centre hospitalier de l'Université de Montréal, Author: Genital Human Papillomavirus (HPV) Infections, Genital Ulcer Disease (GUD), Primary Care and Sexually Transmitted Infections; : **Bruno Turnel**, MD, médecin-conseil, Direction générale de la santé publique, Ministère de la Santé et des Services sociaux du Québec, Author: Epididymitis, Prostatitis, Sexually Transmitted Intestinal and Enteric Infections; **Julie van Schalkwyk**, MD, MSc, FRCSC, Assistant Clinical Professor, Department of Obstetrics and Gynecology, University of British Columbia, Author: Pregnancy; **Tom Wong**, MD, MPH, FRCPC, Director Community Acquired Infections Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada, Author: Chlamydial Infections, Immigrants and Refugees; **Mark H. Yudin**, MD, MSc, FRCSC, Assistant Professor, University of Toronto, Deputy Head, Department of Obstetrics and Gynecology St. Michael's Hospital, Author: Pelvic Inflammatory Disease (PID), Co-Author: Urethritis.

The following people are thanked for volunteering their time as external reviewers for the guideline chapters:

**Robert Brunham**, MD, Director of Medical and Academic Affairs, BC Centre for Disease Control, Director, UBC Centre for Disease Control, Professor of Medicine, Division of Infectious Diseases, University of British Columbia; **Susan Comay**, MD, Medical Director, Sexual Assault Service, BC Women's Hospital; **Curtis Cooper**, MD, FRCPC, University of Ottawa; **Francisco Diaz-Mitoma**, MD, PhD, FRCPC, Professor and Chief, Division of Virology, Children's Hospital of Eastern Ontario, University of Ottawa; **Harold Dion**, MD, CCFP, FCFP, Clinique médicale l'Actuel, Chair of the Board, Quebec College of Family Physicians; **Shelia Dunn**, MD, CCFP(EM), University of Toronto, Sunnybrook and Women's College Health Sciences Centre; **Alex Ferenczy**, MD, Professor of Pathology and Obstetrics & Gynecology, McGill University, Sir Mortimer B. Davis-Jewish General Hospital; **David Fisman**, MD, MPH, Visiting Scholar Center for Health and Wellbeing, Woodrow Wilson School, Princeton University; **Jennifer Geduld**, Manager, HIV/AIDS Surveillance Section, CIDPC, Surveillance and Risk Assessment Division, HIV/AIDS Epidemiology and Surveillance, Public Health Agency of Canada; **Mary Gordon**, MD, City of Ottawa, Public Health, Sexual Health Centre; **Kevin Gough**, MD, FRCPC, MEd, St. Michael's Hospital, University of Toronto; **Andree Gruslin**, MD, FRCS, Post Graduation Program Director, Assistant Professor, MFM Department of Obstetrics and Gynecology, University of Ottawa; **Hunter Handsfield**, MD, Professor of Medicine, University of Washington, Center for AIDS and Sexually Transmitted Diseases; **Sandra Hooper**, RN(EC), MScN, Nurse Practitioner Sexual Health Centre, Ottawa Public Health; **Robbi Howlett**, MAsc, PhD, Manager, Ontario Cervical Screening Program, Division of Preventive Oncology, Cancer Care Ontario; **Gaya Jayaraman**, PhD, MPH, Manager, Hepatitis C and Sexually Transmitted Infections Surveillance Section, Community Acquired Infections Division, Public Health Agency of Canada; **Hugh D Jones**, MD, Dip Ven, STD/AIDS Control, Clinic Physician; **Fadel Kane**, MD, MSc, HIV/AIDS Programs and Policy Division, Public Health Agency of Canada; **Sari Kives**, MD, University of Toronto, St. Michael's Hospital; **Claude Laberge**, MD, Service de lutte contre les ITSS, Direction générale de la santé publique, Ministère de la santé et des services sociaux du Québec; **Gilles Lambert**, MD, Institut national de santé publique du Québec; **Debbie Lindsay**, MD, University of Manitoba, Child Protection Centre, Health Sciences Centre; **Noni MacDonald**, MD, MSc, FRCPC, Dalhousie University, IWK Health Centre; **Louisa MacKenzie**, MD, FRCPC, DTMH, Calgary Refugee Health Program, Margaret Chisholm Resettlement Centre; **Lorette Madore**, RN, DPHN, BN, Supervisor, Clinical Services, Healthy Sexuality and Risk Reduction Program, Ottawa Public Health/Santé publique d'Ottawa; **Cheryl Main**, MD, FRCPC, Assistant Professor Pathology and Molecular Medicine, McMaster University; **Nathalie Mondain**, PhD, Groupe de Recherche Interdisciplinaire en Santé (GRIS), University of Montreal; **Deborah Money**, MD, FRCSC, Associate Professor, University of British Columbia, BC Women's Hospital; Carolyn A Montgomery, MB, STD/AIDS Control, Clinic Physician; **Curtis Nickel**, MD, Professor of Urology, Queen's University; **Gina Ogilvie**, MD, MSc, University

of British Columbia; **Caroline Paquet**, SF, MSc, Professeure, Baccalauréat en pratique sage-femme, Département de Chimie-biologie, Université du Québec à Trois-Rivières; **Gordon Phaneuf**, Director of Strategic Initiatives, Child Welfare League of Canada; **Raphael Saginur**, MD, FRCPC, Ottawa Hospital and University of Ottawa; **John Sellors**, MD, Senior Medical Advisor, Reproductive Health, PATH; **Alberto Severini**, MD, National Microbiology Laboratory, Public Health Agency of Canada; **Stephen Shafran**, MD, FRCPC, Professor and Director, Division of Infectious Diseases, Department of Medicine, University of Alberta; **Rita Shahin**, MD, Toronto Public Health; **Brenna Shearer-Hood**, MSA(HSA), BMR (OT) Cancer Care Manitoba; **Jonathan M. Smith**, CSC Epidemiologist, Tuberculosis Prevention and Control, Community Acquired Infections Division, Public Health Agency of Canada; **Gwen Stephens**, MD, FRCPC, University of British Columbia, Department of Pathology & Laboratory Medicine; **Jill Tinmouth**, MD, PhD, Sunnybrook and Women's Health Sciences Centre, University of Toronto; **Baldwin Toye**, MD, FRCPC, Head, Division of Microbiology, Ottawa Hospital, University of Ottawa; **Sharonie Valin**, MD, CCFP, MHSc, North York General Hospital, Women's College Hospital, Bay Centre for Birth Control; **Heidi Wood**, PhD, Head, Diagnostics, Zoonotic Diseases and Chlamydia Section, National Microbiology Laboratory, Public Health Agency of Canada.

# INTRODUCTION

---

These guidelines were created as a resource for clinical and public health professionals — especially nurses and physicians — for the prevention and management of STIs across a diverse patient population, including neonates, children, adolescents and adults.

Since the release of the *Canadian Guidelines on Sexually Transmitted Infections-2006 Edition*, the Expert Working Group (EWG) for the guidelines which includes STI experts from the fields of medicine, nursing, laboratory, public health and research have volunteered their time and effort as authors and reviewers to maintain updated, evidence-based recommendations for the prevention, diagnosis, treatment and management of STIs in Canada. The content of these current *Canadian Guidelines on Sexually Transmitted Infections* reflects emerging issues and highlight changes in the STI literature since the release of the 2006 guidelines. Guidelines presented in this document reflect the views of the Expert Working Group on Canadian Guidelines for Sexually Transmitted Infections. They should be construed *not* as rules but rather as recommendations.

Canada is a signatory of the WHO global strategy for the prevention and control of sexually transmitted infections 2006-2015 which stipulates that STI prevention and control need to be linked to HIV prevention and sexual and reproductive health programmes and services. The strategy also highlights the need for collaborative efforts at various levels and venues of service delivery for programmes to reach and have the greatest impact on those at risk of STI/HIV and other communicable diseases.

Practitioners should be aware of the emerging research linking STI to preventable chronic diseases in both men and women, such as cancers. Primary care providers are well positioned to offer patient centered STI screening, diagnosis, management and treatment services to a wide range of their patients. Making such a continuum of STI services available, primary care providers have the potential to significantly impact chronic diseases prevention and control in the long-term.

## Future Developments

As within many areas in the health sector, innovation and development are part of the growing body of knowledge and tools used in the prevention, treatment and management of disease and infection. We recommend consulting a variety of mechanisms/sources to maintain and enhance your clinical practice.

## Canadian Guidelines on STI: A New Look

The current binder version of the *Canadian Guidelines on Sexually Transmitted Infections* contains all the updates that were brought to the 2006 Edition and are currently posted on the PHAC website. This new version follows the same section division, chapter order and chapter layout as the previous bound version.

There has been a general index added to the end of the guidelines which highlights which sections key words are found in and then a chapter and page specific index at the end of the sections.

The pagination in the guidelines is now chapter-specific and the section specific index reflects this change. It is anticipated that these changes will facilitate publication of updates for the Guidelines on bi-annual basis, and a two-step index search process (chapter-wise and page-wise) will help readers to navigate the Guidelines easier and faster. The move to a binder format is expected to result in making the Guidelines a live document reflecting the most recent evidence available and making the production and dissemination of updates more time efficient and environmentally friendly.

References for the Canadian Guidelines on Sexually Transmitted Infections are available at: [www.publichealth.gc.ca/sti](http://www.publichealth.gc.ca/sti).

## Need to Strengthen Prevention

In Canada, there are three nationally reportable STIs: chlamydia, gonorrhoea and infectious syphilis. Since 1997, there has been a steady increase in the rates of all three infections. This phenomenon is not unique to Canada; other countries, including the U.S. and the U.K., have reported similar trends.<sup>1,2</sup> Targeted enhanced surveillance and research are required to determine the factors that may be playing a role in these trends. Some of the possible factors may include the following:

- Nucleic acid amplification tests (NAATs) have been introduced.
- Some people may have developed safer-sex burnout.
- There have been innovations in HIV therapy (e.g., highly active antiretroviral therapy), leading to related treatment optimism.
- Youth awareness of risks and knowledge of risk-reduction behaviours remain less than optimal.<sup>3</sup>
- Sex is occurring at an early age, with a high rate of serially monogamous relationships.
- Sex is continuing later in life.
- The transmission risks of STIs associated with sexual activity (vaginal, anal and oral) are not well understood by the public.
- “Party drugs,” such as ecstasy and crystal meth, are being increasingly linked to unsafe sexual behaviours.<sup>4</sup>
- Anonymous partnering venues, such as the Internet, are expanding.

By being aware of trends in STIs, risk factors and affected populations, primary care providers and public health practitioners can be strategically placed to apply relevant and complementary individual and community-based education and patient services.

The prevention and control of STIs cannot be approached with a narrow focus. The appropriate medical management of identified cases of STIs is but one piece of the puzzle. Both primary and secondary prevention activities are paramount to reducing the incidence (newly acquired infections) and prevalence (number of cases) of STIs. Primary prevention aims to prevent exposure by identifying at-risk individuals and performing thorough assessments, patient-centred counselling and education.<sup>5</sup>

Secondary prevention involves reducing the prevalence of STIs through the detection of infections in at-risk populations, counselling, conducting partner notification and treating infected individuals and contacts in a timely manner, thus preventing and/or limiting further spread.<sup>5</sup>

Both the burden of disease and potential complications associated with STIs are relevant and significant considerations for health professionals and decision makers. The presence of an acute infection can increase the risk of co-infection: for example, an ulcer from an infection such as syphilis can significantly increase the risk of acquiring and transmitting an HIV infection. The sequelae for women from untreated infections such as chlamydia and gonorrhoea can include pelvic inflammatory disease, chronic pelvic pain, ectopic pregnancy and infertility. In recent years, there has also been increasing evidence to support the role of persistent human papillomavirus (HPV) infections in cervical dysplasia and carcinoma.

As we strive to attend to the physiological needs of patients, we must also be prepared to attend to their psychological needs as well. Chronic viral STI can have long-standing negative impacts on a patient's psychosocial well-being. The many potential impacts and sequelae of STIs highlight the need for strengthened prevention efforts.

## LEVELS AND QUALITY OF EVIDENCE FOR TREATMENT RECOMMENDATIONS

This updated version contains the same levels of recommendation and quality of evidence indicators for the treatment recommendations as the 2006 Edition. The indicators used reflect a combination of the methodologies from the U.S. Preventive Services Task Force and the Canadian Task Force on Preventive Health Care and have been modified and simplified for use in these guidelines as outlined in Tables 1 and 2.

**Table 1. Levels of recommendation (Modified from Harris RP, et al.<sup>6</sup>)**

<b>Recommendation: A</b>	Strongly recommends that clinicians routinely provide the treatment to eligible patients. <b>Good evidence</b> that the treatment improves important health outcomes and concludes that benefits substantially outweigh harms
<b>Recommendation: B</b>	Recommends that clinicians routinely provide the treatment to eligible patients. At least <b>fair evidence</b> that the treatment improves important health outcomes and concludes that benefits outweigh harms
<b>Recommendation: C</b>	No recommendation for or against routine provision of the treatment. At least <b>fair evidence</b> that the treatment can improve health outcomes but concludes that the balance of the benefits and harms is <b>too close to justify a general recommendation</b>
<b>Recommendation: D</b>	Recommends against routinely providing the treatment to asymptomatic patients. At least <b>fair evidence</b> that the treatment is <b>ineffective</b> or that harms outweigh benefits
<b>Recommendation: I</b>	<b>Evidence is insufficient</b> to recommend for or against routinely providing the treatment. Evidence that the treatment is effective is <b>lacking, of poor quality or conflicting</b> , and the balance of benefits and harms cannot be determined

**Table 2. Quality of evidence (Modified from Harris RP, et al.<sup>6</sup> and Gross PA, et al.<sup>7</sup>)**

<b>I</b>	Evidence from at least one properly randomized, controlled trial
<b>II</b>	Evidence from at least one well-designed clinical trial without randomization, from cohort or case-control analytic studies (preferably from more than one centre), from multiple time-series studies or from dramatic results in uncontrolled experiments
<b>III</b>	Evidence from opinions of respected authorities based on clinical experience, descriptive studies or reports of expert committees

## IMPORTANT CONSIDERATIONS FOR GUIDELINES USERS

The EWG and the PHAC acknowledge that the advice and recommendations set out in this document are based upon the best current available scientific knowledge and medical practices, and they are disseminating this document to clinical and public health professionals for information purposes.

While these guidelines are based on current evidence and clinical practice, the prevention, diagnosis, treatment and management of STIs is an evolving field. The EWG and PHAC will be regularly updating the information and recommendations in the *Guidelines*. Readers are encouraged to consult the PHAC website ([www.publichealth.gc.ca/sti](http://www.publichealth.gc.ca/sti)) for ongoing chapter update(s). In 2008, an electronic database will be developed for the purposes of alerting *Guidelines* users on updates and related issues. If you wish to be notified of the coming updates and provide your feedback or comments on the use of the Guidelines, we ask you to complete the form at the beginning of the document and return it to the Public Health Agency of Canada by email or fax as indicated on the form.

Persons administering or dispensing drugs, vaccines or other products should also be aware of the contents of the individual product monograph(s) for those products, or other similarly approved standards or instructions for use provided by the licensed manufacturer(s). Recommendations for use and other information set out in these guidelines may differ from that set out in product monograph(s) or other similarly approved standards or instructions for use. Manufacturers have sought approval and provided evidence as to the safety and efficacy of their products only when used in accordance with the product monograph(s) or other similarly approved standards or instructions for use.

Practitioners should report adverse drug reactions to the Canadian Adverse Drug Reaction Monitoring Program (CADRMP). For specifications and standards of reporting, consult Health Canada's CADRMP guidelines.

**While this document addresses key issues related to the prevention, diagnosis, treatment and management of the most common STIs, it is beyond the scope of these guidelines to provide comprehensive recommendations for the treatment and management of HIV and viral hepatitis C. When confronted with these infections, either as a primary infection or a co-infection, we suggest that you refer to alternate resources (see below for suggestions), including colleagues experienced in the area.**

- Strader DB, Wright T, Thomas DL, Seeff LB. AASLD practice guideline: diagnosis, management, and treatment of hepatitis C. *Hepatology* 2004;39:1147–1171.
- U.S. Department of Health and Human Services, Panel on Clinical Practices for Treatment of HIV Infection. *Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents*. Available at: <http://www.aidsinfo.nih.gov/Guidelines/GuidelineDetail.aspx?MenuItem=Guidelines&Search=Off&GuidelineID=7&ClassID=1> (Accessed December 19, 2007)